



## King's Research Portal

DOI:

[10.1016/j.coph.2018.08.004](https://doi.org/10.1016/j.coph.2018.08.004)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Simpson, S., Smith, L., & Bowe, J. (2018). Placental peptides regulating islet adaptation to pregnancy: clinical potential in gestational diabetes mellitus: • Endocrine & Metabolic Diseases • Gastrointestinal. *Current Opinion in Pharmacology*, 43, 59-65. <https://doi.org/10.1016/j.coph.2018.08.004>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Placental peptides regulating islet adaptation to pregnancy: clinical potential in gestational diabetes mellitus

Sian Simpson, Lorna Smith and James Bowe



Pregnancy involves a progressive increase in insulin resistance and the  $\beta$ -cells must adapt to compensate and prevent gestational diabetes (GDM). In this review we discuss the evidence for placental peptides, including placental lactogen, hepatocyte growth factor, adiponectin and leptin, playing a role in the islet adaptation to pregnancy. The difficulties of translating data from rodent models into human pregnancy are covered and we summarise studies investigating associations between serum placental peptides and GDM risk. In conclusion, current data support important roles for placental peptides interacting to support  $\beta$ -cells during pregnancy, however mechanisms involved in humans are unclear. Further work in humans is required, but placental peptides have clinical potential from both a diagnostic and therapeutic perspective.

## Address

Department of Diabetes, School of Life Course Sciences, King's College London, Guy's Campus, London SE1 1UL, UK

Corresponding author: Bowe, James ([james.bowe@kcl.ac.uk](mailto:james.bowe@kcl.ac.uk))

Current Opinion in Pharmacology 2018, 43:59–65

This review comes from a themed issue on **Endocrine and metabolic diseases**

Edited by **James Bowe** and **Shanta Persaud**

<https://doi.org/10.1016/j.coph.2018.08.004>

1471-4892/© 2018 Published by Elsevier Ltd.

## Introduction

During a healthy pregnancy insulin sensitivity falls with gestational age. This physiological change prioritises the delivery of glucose across the placenta for fetal development. However, this poses a dilemma for the maternal metabolism to balance providing for the energy requirements of the growing fetus, while simultaneously maintaining maternal glucose homeostasis. In a healthy mother the insulin resistance is countered by adaptive changes in pancreatic islets to allow increased insulin secretion, including an increased  $\beta$ -cell mass. Failure of the insulin-releasing  $\beta$ -cells to sufficiently adapt and compensate for the increased metabolic demand in pregnancy leads to development

of glucose intolerance, hyperglycaemia and gestational diabetes mellitus (GDM).

## Gestational diabetes

The rapid worldwide increase in the prevalence of type 2 diabetes mellitus (T2DM) is well-documented, but it is less appreciated that the incidence of GDM is also rapidly rising in parallel with the T2DM pandemic. It is currently estimated worldwide that 21.3 million pregnancies per year (16.2% of total pregnancies) are affected by some form of hyperglycaemia, with 86.4% of those cases due to GDM (IDF Diabetes Atlas: <https://www.idf.org/e-library/epidemiology-research/diabetes-atlas/134-idf-diabetes-atlas-8th-edition.html>). Maternal GDM has consequences for the mother and developing fetus, including placental insufficiency because of preeclampsia, macrosomia (birth weight >90th centile), birth injury and neonatal hyperinsulinism and hypoglycaemia. Epidemiological and experimental studies have also demonstrated that the GDM intra-uterine environment can increase susceptibility of the offspring to later life T2DM and cardiovascular disease, whilst in mothers GDM is associated with the subsequent development of T2DM.

At present GDM is diagnosed through routine oral glucose tolerance testing at 24–28 weeks of gestation, once impaired glucose tolerance has developed. Therapeutically GDM is primarily treated with either insulin injections or metformin, though the use of insulin analogs which do not cross the placenta have also been suggested [1]. Given the range of acute and potentially chronic consequences of GDM for both mother and child, there is currently great interest in developing diagnostic tools for predicting high GDM risk earlier in pregnancy, thus allowing time for intervention. Additionally, the identification of additional therapeutic targets is of keen interest for investigation.

## Placental peptides

GDM is primarily the result of insufficient islet adaptation and an inability to increase insulin secretory capacity to compensate for increased insulin resistance. In normal pregnancy  $\beta$ -cells adapt in several respects to increase functionality. In rodent models these mechanisms include increased insulin synthesis and release, increased glucose responsiveness, increased cell–cell communication, hypertrophy, increased proliferation and reduced apoptosis, resulting in an overall increase in  $\beta$ -cell mass.

The  $\beta$ -cell adaptation in human pregnancy is more controversial, particularly concerning  $\beta$ -cell proliferation, but broadly speaking a similar pattern is observed with increased insulin release and increased  $\beta$ -cell mass. Thus, understanding the signals and mechanisms that regulate the islet adaptation to pregnancy and the reasons for these mechanisms failing in GDM warrants further investigation.

The placenta plays many essential roles during pregnancy, including supporting and protecting the fetus, and the supply of nutrients and gas exchange. The placenta also acts as an essential endocrine organ, producing and releasing hormones and mediators into the maternal circulation to maintain pregnancy. Increasing evidence suggests that several placental hormones play a role in communicating with maternal  $\beta$ -cells to regulate the islet adaptation necessary for a healthy pregnancy.

The effects of steroid hormones released from the placenta, including oestrogens, progestogens and glucocorticoids, on glucose homeostasis are complex and still controversial. Oestrogens have been implicated in the islet adaptation to pregnancy, through both direct and indirect protective effects on  $\beta$ -cells [2]. Effects of progestogens appear to vary with concentration and the presence of other hormones [2], and further work is required to understand their role. The placenta also releases a much wider range of peptide hormones into the maternal circulation that may influence the islet adaptation to pregnancy. Despite  $\beta$ -cells expressing cognate receptors for many of these ligands, only a few have been investigated for possible effects during pregnancy. Thus, this review will summarise the placental peptides currently thought to play a role in the islet adaptation to pregnancy and discuss whether these peptides may represent useful biomarkers for early determination of GDM risk.

### Lactogenic hormones

The lactogenic hormones, prolactin (PRL) and placental lactogen (PL), are the most extensively studied hormones involved in the islet adaptation to pregnancy. During early pregnancy PRL is released from the maternal anterior pituitary, but following placentation maternal circulating levels of PRL fall and PL released from the placenta becomes the dominant lactogenic signal during peak  $\beta$ -cell adaptation. Both PRL and PL exert effects through the prolactin receptor (PRLR), which is expressed specifically on the  $\beta$ -cell of rodent islets [3].

The role of PL in the islet adaptation to pregnancy was initially implicated largely through studies demonstrating that lactogens increased glucose-induced insulin secretion,  $\beta$ -cell proliferation and survival in isolated rodent

islets [4–6]. Global homozygous *Prhr* knockout mice exhibit reproductive dysfunction [7], so are unsuitable for pregnant studies. However, heterozygous *Prhr*+/- mice are fertile and have impaired glucose tolerance and reduced  $\beta$ -cell mass during pregnancy [8], but also adipose tissue effects that influence glucose homeostasis. The recently established  $\beta$ -cell specific *Prhr* knockout mouse ( $\beta$ PRLR-KO) provides the best evidence for the role of lactogenic hormones in  $\beta$ -cell adaptation. Although  $\beta$ PRLR-KO mice have normal glucose tolerance outside of pregnancy, they become progressively glucose intolerant during gestation, with corresponding failure of  $\beta$ -cell proliferation [9\*\*].

The intracellular mechanisms activated by lactogenic hormones *in vitro* closely reflect pregnant changes, including glucokinase upregulation [10] and pro-proliferative and anti-apoptotic signalling pathways [11,12]. The lactogenic hormones also stimulate  $\beta$ -cell production of serotonin during pregnancy [13], which appears to play a critical local role in regulating  $\beta$ -cell mass [14], mediating the effects of prolactin (Table 1).

Given the role for lactogenic hormones in rodent pregnancy, there is ongoing research to determine whether similar mechanisms are also relevant in human  $\beta$ -cells. Human PLs are derived from duplications of the *hGH* gene, whilst rodent PLs are evolved from the prolactin gene [15], and the *Prhr* gene is significantly enriched in mouse  $\beta$ -cells compared to human, perhaps indicating a lesser role in human  $\beta$ -cell adaptation [16,17]. Treatment of human islets with lactogenic hormones potentiates glucose-stimulated insulin secretion [18], but the effects on  $\beta$ -cell proliferation are more controversial. Increased proliferation, as assessed by BrdU incorporation, has been reported in human islets in response to PL and PRL [18], but more recent studies have been unable to replicate this effect [19].

It is also unclear whether serum PRL or PL associates with GDM pathophysiology. Several studies have investigated possible links, but the majority found no significant correlation between maternal PRL or PL and GDM [20], although counter-intuitively an association between high maternal prolactin and reduced glucose tolerance has recently been reported [21\*\*]. Despite the apparent lack of association to GDM risk, low PRL or PL levels have been associated with reduced glucose tolerance post-partum and an increased risk of subsequent pre-diabetes/diabetes [22\*], suggesting that the lactogenic hormones do play some role in human pregnancy.

### Hepatocyte growth factor

Hepatocyte growth factor (HGF), acting via the c-Met receptor, has also been implicated in the islet adaptation to pregnancy. Normally HGF is released from endothelial cells in the islet vasculature to exert local effects on the  $\beta$ -cells [23], however during pregnancy high levels of

**Table 1****Effects of placental peptides investigated in rodent and human islets that may potentially play a role in the islet adaptation to pregnancy and the development of GDM**

Placental signal	Effects on rodent islets	Effects on human islets	Correlation with GDM
Placental lactogen	Increased glucose stimulated insulin secretion, increased proliferation and pro-survival effects, upregulation of islet serotonin	Controversial, but some studies report increased glucose-stimulated insulin secretion and proliferation	No significant correlation/high PL associated with increased GDM risk
HGF	Pro-proliferative and anti-apoptotic effects	Anti-apoptotic effects in human fetal islets and transplanted human islet grafts	High HGF associated with increased GDM risk in obese women
Leptin	Reduced glucose-induced insulin secretion and potentially $\beta$ -cell atrophy	Reduced glucose-induced insulin secretion, effects on $\beta$ -cell mass unknown	High leptin associated with increased GDM risk
Adiponectin	Increased glucose-stimulated insulin secretion and $\beta$ -cell mass	No effect on insulin secretion, effects on $\beta$ -cell mass unknown	Low adiponectin associated with GDM risk
Kisspeptin	Increased glucose-induced insulin secretion, effects on $\beta$ -cell mass unknown	Increased glucose-induced insulin secretion, effects on $\beta$ -cell mass unknown	Low kisspeptin associated with GDM risk

circulating HGF are also released from the placenta [24]. Loss of pancreatic c-Met signalling has no significant effect on glucose homeostasis outside of pregnancy [25], but results in reduced  $\beta$ -cell proliferation and increased apoptosis during pregnancy, with consequently reduced plasma insulin and impaired glucose tolerance [26]. Interestingly, pancreatic c-Met knockout mice also lack the increase in islet *Prlr* levels seen in normal pregnancy, suggesting that HGF may represent an upstream factor regulating PRLR signalling [26].

Whilst few studies have examined effects of HGF on human islet function in relation to pregnancy, HGF does stimulate proliferation in human fetal  $\beta$ -cells [27] and improves graft survival of human islets through anti-apoptotic effects in islet transplantation models [28,29]. Despite these effects of HGF in human  $\beta$ -cells, there does not appear to be any correlation between serum HGF levels and GDM progression across the whole population [30]. However, obese women generally have significantly higher placental HGF levels when compared to control women [31], and specifically among obese women high serum HGF is associated with a 4.5-fold higher risk of developing GDM [30].

Given the proliferative and pro-survival effects of HGF on  $\beta$ -cells it is surprising that high HGF appears to be deleterious. The relative contributions of placental and endothelial HGF on  $\beta$ -cell adaptation are unclear. As placental HGF rises through gestation, islet endothelial cells also proliferate, increasing local HGF. This has been identified as a possible mechanism for coordinating vascular and  $\beta$ -cell function in pregnancy [23]. Given that raised placental HGF associates with increased GDM risk in obese women, local endothelial cell-derived HGF could be the primary driver of  $\beta$ -cell responses. High

HGF does associate with GDM risk in obese women and as such may be a useful biomarker for GDM generally, but it is unlikely to be an indicator of  $\beta$ -cell adaptation.

### Adipokines

The adipokines are signalling molecules secreted by the adipocytes that regulate processes including atherosclerosis, inflammation, appetite and glucose homeostasis. The adipokines are present at increasing levels as adipose tissue expands in obesity and several adipokines, including leptin, adiponectin, visfatin, chemerin and omentin-1 have been implicated in the development of T2DM [32]. Leptin and adiponectin are also released at high levels from the placenta into the maternal circulation and, although they are more commonly linked to effects on food intake or insulin sensitivity, they have additional direct effects on  $\beta$ -cells.

Adiponectin has primarily been studied as a regulator of insulin sensitivity, but both human and rodent  $\beta$ -cells express the adiponectin receptors AdipoR1 and AdipoR2 [33]. Adiponectin has been shown to stimulate insulin secretion from mouse islets both *in vitro* and *in vivo* [34,35], although the same effect is not observed in human islets [36]. Pregnant adiponectin knockout mice have reduced  $\beta$ -cell mass and insulin deficiency, though despite this they are able to maintain normal plasma insulin levels [37<sup>••</sup>]. However, these mice also display increased triglyceride production, hyperlipidaemia and increased hepatic glucose, which makes assessing the relative importance of  $\beta$ -cell-specific adiponectin signalling on glucose tolerance difficult [37<sup>••</sup>].

Clinical studies have found decreased serum adiponectin in women with GDM compared to women with healthy pregnancies [38–41]. Placental adiponectin expression is also downregulated in GDM, suggesting that reduced

serum adiponectin in GDM is at least partly due to low placental, as opposed to adipocyte, secretion [42]. Whilst adiponectin clearly regulates  $\beta$ -cell adaptation to pregnancy, it also has a well-established role in increasing insulin sensitivity. Whether the link between circulating adiponectin and GDM risk is the consequence of one or both of these mechanisms is currently unclear, but first trimester adiponectin has been suggested as a potentially useful biomarker for predicting GDM [43\*\*].

The hypothalamic effects of leptin to regulate food intake, with subsequent effects on adiposity, insulin resistance and glucose tolerance are well-established. However, mouse and human  $\beta$ -cells also express several isoforms of the leptin receptor [44]. Leptin reduces glucose-induced insulin secretion in isolated mouse islets [45,46] when administered to mice *in vivo* [47] and in isolated human islets [47,48]. Mice heterozygous for the leptin receptor (*db/+*) develop GDM during pregnancy, with associated glucose intolerance and macrosomia, though they also exhibit increased weight gain and adiposity [49]. Whether the impaired glucose tolerance is due to direct  $\beta$ -cell effects or secondary to hypothalamic effects of leptin is unknown. Mice with  $\beta$ -cell-specific leptin receptor knockout show mildly improved glucose tolerance, enhanced glucose-stimulated insulin secretion and increased  $\beta$ -cell mass through hypertrophy [50,51], but have not yet been studied in pregnancy.

There is a well-established association between raised serum leptin and increased GDM risk [52\*,53], which has resulted in leptin having also been proposed as a first trimester biomarker for establishing GDM risk [52\*,54]. Similar to adiponectin, circulating leptin in pregnancy may be derived from either the placenta or the adipocytes, but leptin expression is increased in GDM placenta [55–57]. As with adiponectin, the relative importance of the  $\beta$ -cell and the hypothalamic effects of leptin on GDM risk is unclear.

### Novel signals

The signals described above are all placental derived hormones whose potential role in mediating the islet adaptation to pregnancy has been established to some extent. However, it is worth noting that the placenta produces and secretes a very wide range of different peptide hormones into the maternal circulation. G-protein coupled receptors (GPCRs) represent the most abundant family of cell surface receptors and mouse and human islets express 279 and 293 different GPCRs respectively [58]. Additionally the mouse placenta expresses mRNA for 79 different endogenous GPCR ligands, the majority of which have at least one cognate receptor on the islets [59\*\*]. As yet unidentified signals may also play a significant role in the communication between placenta and  $\beta$ -cells.

Kisspeptin represents one placental GPCR ligand that potentially plays a role in the  $\beta$ -cell adaptive response to pregnancy. Under normal physiological conditions kisspeptin is primarily a hypothalamic signal involved in regulating reproductive function, but during pregnancy serum levels increase several-thousand fold due to placental release [60]. Both mouse and human islets express the kisspeptin receptor, GPR54 [61], and kisspeptin stimulates glucose-stimulated insulin secretion in isolated mouse and human islets [62,63]. Unfortunately, due to the role of kisspeptin in reproductive function, global GPR54 knockout mice are infertile and unsuitable for pregnancy studies [64]. However, upcoming studies in  $\beta$ -cell specific GPR54 knockout mice show impaired glucose tolerance and reduced glucose-stimulated insulin secretion during pregnancy (T. Hill, abstract in Diabetic Medicine 2017, 34(S1): 27–28). Furthermore, clinical studies have shown that serum kisspeptin is significantly lower in GDM [65] and can be safely administered in humans to stimulate glucose-stimulated insulin secretion [66\*\*]. More work is required to determine whether kisspeptin represents a potentially important signal to add to the established factors described above.

### Conclusions

There is a growing body of evidence linking placental peptides to rodent  $\beta$ -cell adaptation to pregnancy. However, the major challenge going forward is understanding how these mechanisms translate to human pregnancy. How human  $\beta$ -cells adapt to pregnancy is still poorly understood, particularly in terms of  $\beta$ -cell mass. Post-mortem studies indicate that  $\beta$ -cell mass increases during human pregnancy, but the mechanisms remain controversial. The limited data suggests  $\beta$ -cell proliferation does not increase in human pregnancy, as assessed by Ki67 staining, and neogenesis is more important in humans [67]. These conclusions are based on autopsy samples from different time-points when there may not have been active proliferation, but they demonstrate the difficulties in translating results from rodent models into humans. The core principles associated with  $\beta$ -cell adaptation in rodents also appear to be conserved to at least some extent in human pregnancy, though a clearer understanding of this is key for future advances.

Understanding the signals involved in regulating the human  $\beta$ -cell response to pregnancy is of great interest for both potential diagnostic and therapeutic benefits. There are ongoing efforts to develop a panel of biomarkers to identify mothers at high risk of GDM before glucose intolerance develops. Most currently suggested biomarkers are steroid hormones and metabolites [52\*], but peptide hormones may be a useful addition. Adiponectin and leptin would appear the most promising candidates to screen for and identify GDM risk, whilst kisspeptin represents a novel candidate that may also be promising with further research.



In terms of therapeutic potential far more work is required. The PL, HGF and the adipokines have off-target effects and may consequently not be useful as therapies. Perhaps the most promising therapeutic candidates are novel placental GPCR ligands. One-third of all prescription drugs in use target GPCRs, including GLP-1 receptor agonists for T2DM, placental GPCR ligands could include as yet unidentified therapeutics. Indeed kisspeptin has recently been shown to safely stimulate glucose-induced insulin secretion in humans [66•] and is currently the subject of clinical trials (NIH Clinical Trials: <https://clinicaltrials.gov/ct2/show/study/NCT02953834>) which may be of future use in pregnancy.

In summary, key placental peptides have been identified to play roles in regulating the islet adaptation to pregnancy in rodent models. Some signals may be of diagnostic use in biomarker screening for GDM risk. However, our understanding of  $\beta$ -cell adaptation to pregnancy in humans is still at a relatively early stage and further studies are required to identify potentially safe and effective therapeutics.

## Conflict of interest

Nothing declared.

## Acknowledgements

The authors and their work in this area is supported by grant funding from the Wellcome Trust (grant number 108064/Z/15/Z), the BBSRC (grant number BB/N00616X/1) and the Diabetes Research and Wellness Foundation.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as

- of special interest
- of outstanding interest

1. Hedrington MS, Davis SN: **The care of pregestational and gestational diabetes and drug metabolism considerations.** *Expert Opin Drug Metab Toxicol* 2017, **13**:1029-1038.
2. Mauvais-Jarvis F: **Role of sex steroids in beta cell function, growth, and survival.** *Trends Endocrinol Metab* 2016, **27**:844-855.
3. Brelje TC, Svensson AM, Stout LE, Bhagroo NV, Sorenson RL: **An immunohistochemical approach to monitor the prolactin-induced activation of the JAK2/STAT5 pathway in pancreatic islets of Langerhans.** *J Histochem Cytochem* 2002, **50**:365-383.
4. Parsons JA, Brelje TC, Sorenson RL: **Adaptation of islets of Langerhans to pregnancy: increased islet cell proliferation and insulin secretion correlates with the onset of placental lactogen secretion.** *Endocrinology* 1992, **130**:1459-1466.
5. Sorenson RL, Brelje TC: **Adaptation of islets of Langerhans to pregnancy: beta-cell growth, enhanced insulin secretion and the role of lactogenic hormones.** *Horm Metab Res* 1997, **29**:301-307.
6. Sorenson RL, Brelje TC, Roth C: **Effects of steroid and lactogenic hormones on islets of Langerhans: a new hypothesis for the role of pregnancy steroids in the adaptation of islets to pregnancy.** *Endocrinology* 1993, **133**:2227-2234.
7. Ormandy CJ, Camus A, Barra J, Damotte D, Lucas B, Buteau H, Edery M, Brousse N, Babinet C, Binart N *et al.*: **Null mutation of the prolactin receptor gene produces multiple reproductive defects in the mouse.** *Genes Dev* 1997, **11**:167-178.
8. Huang C, Snider F, Cross JC: **Prolactin receptor is required for normal glucose homeostasis and modulation of beta-cell mass during pregnancy.** *Endocrinology* 2009, **150**:1618-1626.
9. Banerjee RR, Cyphert HA, Walker EM, Chakravarthy H, Peiris H, Gu X, Liu Y, Conrad E, Goodrich L, Stein RW *et al.*: **Gestational diabetes mellitus from inactivation of prolactin receptor and MafB in islet beta-cells.** *Diabetes* 2016, **65**:2331-2341.
- This study generates a  $\beta$ -cell specific PRLR knockout mouse and shows that loss of  $\beta$ -cell lactogen signalling results in gestational diabetes.
10. Weinhaus AJ, Stout LE, Sorenson RL: **Glucokinase, hexokinase, glucose transporter 2, and glucose metabolism in islets during pregnancy and prolactin-treated islets in vitro: mechanisms for long term up-regulation of islets.** *Endocrinology* 1996, **137**:1640-1649.
11. Banerjee RR: **Piecing together the puzzle of pancreatic islet adaptation in pregnancy.** *Ann N Y Acad Sci* 2018, **1411**:120-139.
12. Rieck S, Kaestner KH: **Expansion of beta-cell mass in response to pregnancy.** *Trends Endocrinol Metab* 2010, **21**:151-158.
13. Schraenen A, Lemaire KdeFG, Hendrickx N, Granvik M, Van LL, Mallet J, Vodjdani G, Gilon P, Binart N *et al.*: **Placental lactogens induce serotonin biosynthesis in a subset of mouse beta cells during pregnancy.** *Diabetologia* 2010, **53**:2589-2599.
14. Kim H, Toyofuku Y, Lynn FC, Chak E, Uchida T, Mizukami H, Fujitani Y, Kawamori R, Miyatsuka T, Kosaka Y *et al.*: **Serotonin regulates pancreatic beta cell mass during pregnancy.** *Nat Med* 2010, **16**:804-808.
15. Haig D: **Placental growth hormone-related proteins and prolactin-related proteins.** *Placenta* 2008, **29**:S36-S41.
16. Benner C, van der Meulen T, Caceres E, Tigyi K, Donaldson CJ, Huising MO: **The transcriptional landscape of mouse beta cells compared to human beta cells reveals notable species differences in long non-coding RNA and protein-coding gene expression.** *BMC Genomics* 2014, **15**:620.
17. Xin Y, Kim J, Okamoto H, Ni M, Wei Y, Adler C, Murphy AJ, Yancopoulos GD, Lin C, Gromada J, Kosaka Y *et al.*: **RNA sequencing of single human islet cells reveals type 2 diabetes genes.** *Cell Metab* 2016, **24**:608-615.
18. Brelje TC, Scharp DW, Lacy PE, Ogren L, Talamantes F, Robertson M, Friesen HG, Sorenson RL: **Effect of homologous placental lactogens, prolactins, and growth hormones on islet B-cell division and insulin secretion in rat, mouse, and human islets: implication for placental lactogen regulation of islet function during pregnancy.** *Endocrinology* 1993, **132**:879-887.
19. Chen H, Kleinberger JW, Takane KK, Salim F, Fiaschi-Taesch N, Pappas K, Parsons R, Jiang J, Zhang Y, Liu H *et al.*: **Augmented Stat5 signaling bypasses multiple impediments to lactogen-mediated proliferation in human beta-cells.** *Diabetes* 2015, **64**:3784-3797.
20. Retnakaran R, Ye C, Kramer CK, Connelly PW, Hanley AJ, Sermer M, Zinman B: **Evaluation of circulating determinants of beta-cell function in women with and without gestational diabetes.** *J Clin Endocrinol Metab* 2016, **101**:2683-2691.
21. Ekinci EI, Torkamani N, Ramchand SK, Churilov L, Sikaris KA, Lu ZX, Houlihan CA: **Higher maternal serum prolactin levels are associated with reduced glucose tolerance during pregnancy.** *J Diabetes Investig* 2017, **8**:697-700.
- Previous clinical studies have generally found no association between serum lactogenic hormones, but this recent study suggests that high maternal lactogens may associate with GDM risk.
22. Retnakaran R, Ye C, Kramer CK, Connelly PW, Hanley AJ, Sermer M, Zinman B: **maternal serum prolactin and prediction of postpartum beta-cell function and risk of prediabetes/diabetes.** *Diabetes Care* 2016, **39**:1250-1258.
- This clinical study finds that whilst the lactogenic hormones do not appear to associate with GDM risk, there is an intriguing correlation between low serum PRL/PL and post-partum beta-cell function.
23. Johansson M, Mattsson G, Andersson A, Jansson L, Carlsson PO: **Islet endothelial cells and pancreatic beta-cell proliferation: studies in vitro and during pregnancy in adult rats.** *Endocrinology* 2006, **147**:2315-2324.

24. Horibe N, Okamoto T, Itakura A, Nakanishi T, Suzuki T, Kazeto S, Tomoda Y: **Levels of hepatocyte growth factor in maternal serum and amniotic fluid.** *Am J Obstet Gynecol* 1995, **173**:937-942.
25. Mellado-Gil J, Rosa TC, Demirci C, Gonzalez-Pertusa JA, Velazquez-Garcia S, Ernst S, Valle S, Vasavada RC, Stewart AF, Alonso LC *et al.*: **Disruption of hepatocyte growth factor/c-Met signaling enhances pancreatic beta-cell death and accelerates the onset of diabetes.** *Diabetes* 2011, **60**:525-536.
26. Demirci C, Ernst S, Alvarez-Perez JC, Rosa T, Valle S, Shridhar V, Casinelli GP, Alonso LC, Vasavada RC, Garcia-Ocana A: **Loss of HGF/c-Met signaling in pancreatic beta-cells leads to incomplete maternal beta-cell adaptation and gestational diabetes mellitus.** *Diabetes* 2012, **61**:1143-1152.
27. Beattie GM, Rubin JS, Mally MI, Otonkoski T, Hayek A: **Regulation of proliferation and differentiation of human fetal pancreatic islet cells by extracellular matrix, hepatocyte growth factor, and cell-cell contact.** *Diabetes* 1996, **45**:1223-1228.
28. Fiaschi-Taesch N, Stewart AF, Garcia-Ocana A: **Improving islet transplantation by gene delivery of hepatocyte growth factor (HGF) and its downstream target, protein kinase B (PKB)/Akt.** *Cell Biochem Biophys* 2007, **48**:191-199.
29. Yeung TY, Seeberger KL, Kin T, Adesida A, Jomha N, Shapiro AM, Korbitt GS: **Human mesenchymal stem cells protect human islets from pro-inflammatory cytokines.** *PLOS ONE* 2012, **7**: e38189.
30. Dishi M, Hevner K, Qiu C, Fida NG, Abetew DF, Williams MA, Enquobahrie DA: **Early pregnancy maternal hepatocyte growth factor and risk of gestational diabetes.** *Br J Med Med Res* 2015, **9**.
31. Visiedo F, Bugatto F, Carrasco-Fernandez C, Saez-Benito A, Mateos RM, Cozar-Castellano I, Bartha JL, Perdomo G: **Hepatocyte growth factor is elevated in amniotic fluid from obese women and regulates placental glucose and fatty acid metabolism.** *Placenta* 2015, **36**:381-388.
32. Rabe K, Lehrke M, Parhofer KG, Broedl UC: **Adipokines and insulin resistance.** *Mol Med* 2008, **14**:741-751.
33. Kharroubi I, Rasschaert J, Eizirik DL, Chop M: **Expression of adiponectin receptors in pancreatic beta cells.** *Biochem Biophys Res Commun* 2003, **312**:1118-1122.
34. Cantley J: **The control of insulin secretion by adipokines: current evidence for adipocyte-beta cell endocrine signalling in metabolic homeostasis.** *Mamm Genome* 2014, **25**:442-454.
35. Okamoto M, Ohara-Imaizumi M, Kubota N, Hashimoto S, Eto K, Kanno T, Kubota T, Wakui M, Nagai R, Noda M *et al.*: **Adiponectin induces insulin secretion in vitro and in vivo at a low glucose concentration.** *Diabetologia* 2008, **51**:827-835.
36. Staiger K, Stefan N, Staiger H, Brendel MD, Brandhorst D, Bretzel RG, Machicao F, Kellerer M, Stumvoll M, Fritsche A *et al.*: **Adiponectin is functionally active in human islets but does not affect insulin secretory function or beta-cell lipapoptosis.** *J Clin Endocrinol Metab* 2005, **90**:6707-6713.
37. Qiao L, Watzet JS, Lee S, Nguyen A, Schaack J, Hay WW Jr, Shao J: **Adiponectin deficiency impairs maternal metabolic adaptation to pregnancy in mice.** *Diabetes* 2017, **66**:1126-1135.
- This recent study investigates the effects of global adiponectin deficiency on glucose homeostasis in pregnant mice, including effects on the islets.
38. Bhograj A, Suryanarayana KM, Nayak A, Murthy NS, Dharmalingam M, Kalra P: **Serum adiponectin levels in gestational diabetes mellitus.** *Indian J Endocrinol Metab* 2016, **20**:752-755.
39. Pala HG, Ozalp Y, Yener AS, Gerçeklioglu G, Uysal S, Onvural A: **Adiponectin levels in gestational diabetes mellitus and in pregnant women without glucose intolerance.** *Adv Clin Exp Med* 2015, **24**:85-92.
40. Ramirez VI, Miller E, Meireles CL, Gelfond J, Krummel DA, Powell TL: **Adiponectin and IGFBP-1 in the development of gestational diabetes in obese mothers.** *BMJ Open Diabetes Res Care* 2014, **2**:e000010.
41. Retnakaran R, Hanley AJ, Raif N, Connelly PW, Sermer M, Zinman B: **Reduced adiponectin concentration in women with gestational diabetes: a potential factor in progression to type 2 diabetes.** *Diabetes Care* 2004, **27**:799-800.
42. Chen J, Tan B, Karteris E, Zervou S, Digby J, Hillhouse EW, Vavish M, Rande HS: **Secretion of adiponectin by human placenta: differential modulation of adiponectin and its receptors by cytokines.** *Diabetologia* 2006, **49**:1292-1302.
43. Thagaard IN, Krebs L, Holm JC, Lange T, Larsen T, Christiansen M: **Adiponectin and leptin as first trimester markers for gestational diabetes mellitus: a cohort study.** *Clin Chem Lab Med* 2017, **55**:1805-1812.
- This clinical study investigates the potential of adiponectin and leptin as early predictors of GDM, finding that both may be potentially useful biomarkers but particularly adiponectin.
44. Marroqui L, Gonzalez A, Neco P, Caballero-Garrido E, Vieira E, Ripoll C, Nadal A, Quesada I: **Role of leptin in the pancreatic beta-cell: effects and signaling pathways.** *J Mol Endocrinol* 2012, **49**:R9-R17.
45. Kieffer TJ, Heller RS, Leech CA, Holz GG, Habener JF: **Leptin suppression of insulin secretion by the activation of ATP-sensitive K<sup>+</sup> channels in pancreatic beta-cells.** *Diabetes* 1997, **46**:1087-1093.
46. Zhao AZ, Bornfeldt KE, Beavo JA: **Leptin inhibits insulin secretion by activation of phosphodiesterase 3B.** *J Clin Invest* 1998, **102**:869-873.
47. Kulkarni RN, Wang ZL, Wang RM, Hurley JD, Smith DM, Ghate MA, Withers DJ, Gardiner JV, Bailey CJ, Bloom SR: **Leptin rapidly suppresses insulin release from insulinoma cells, rat and human islets and, in vivo, in mice.** *J Clin Invest* 1997, **100**:2729-2736.
48. Lupi R, Marchetti P, Maffei M, Del Guerra S, Benzi L, Marselli L, Bertacca A, Navalesi R: **Effects of acute or prolonged exposure to human leptin on isolated human islet function.** *Biochem Biophys Res Commun* 1999, **256**:637-641.
49. Yamashita H, Shao J, Ishizuka T, Klepcyk PJ, Muhlenkamp P, Qiao L, Hoggard N, Friedman JE: **Leptin administration prevents spontaneous gestational diabetes in heterozygous Lepr(db/+) mice: effects on placental leptin and fetal growth.** *Endocrinology* 2001, **142**:2888-2897.
50. Morioka T, Asilmaz E, Hu J, Dishinger JF, Kurpad AJ, Elias CF, Li H, Elmquist JK, Kennedy RT, Kulkarni RN: **Disruption of leptin receptor expression in the pancreas directly affects beta cell growth and function in mice.** *J Clin Invest* 2007, **117**:2860-2868.
51. Soedling H, Hodson DJ, Adrianssens AE, Gribble FM, Reimann F, Trapp S, Rutter GA: **Limited impact on glucose homeostasis of leptin receptor deletion from insulin- or proglucagon-expressing cells.** *Mol Metab* 2015, **4**:619-630.
52. Powe CE: **Early pregnancy biochemical predictors of gestational diabetes mellitus.** *Curr Diab Rep* 2017, **17**:12.
- This recent paper covers the different biochemical markers that are currently thought to be most potentially useful for GDM screening.
53. Tsiotra PC, Halvatsiotis P, Patsouras K, Maratou E, Salamalekis G, Raptis SA, Dimitriadis G, Boutati E: **Circulating adipokines and mRNA expression in adipose tissue and the placenta in women with gestational diabetes mellitus.** *Peptides* 2018, **101**:157-166.
54. Iciek R, Wender-Ozegowska E, Zawiejska A, Mikolajczak P, Mrozikiewicz PM, Pietryga M, Brazert J: **Placental leptin and its receptor genes expression in pregnancies complicated by type 1 diabetes.** *J Physiol Pharmacol* 2013, **64**:579-585.
55. Perez-Perez A, Guadix P, Maymo J, Duenas JL, Varone C, Fernandez-Sanchez M, Sanchez-Margalet V: **Insulin and leptin signaling in placenta from gestational diabetic subjects.** *Horm Metab Res* 2016, **48**:62-69.
56. Perez-Perez A, Maymo JL, Gambino YP, Guadix P, Duenas JL, Varone CL, Sanchez-Margalet V: **Activated translation signaling in placenta from pregnant women with gestational diabetes mellitus: possible role of leptin.** *Horm Metab Res* 2013, **45**:436-442.

57. Lepercq J, Cauzac M, Lahlou N, Timsit J, Girard J, Auwerx J, Hauguel-de Mouzon S: **Overexpression of placental leptin in diabetic pregnancy: a critical role for insulin.** *Diabetes* 1998, **47**:847-850.
  58. Amisten S, Atanes P, Hawkes R, Ruz-Maldonado I, Liu B, Parandeh F, Zhao M, Huang GC, Salehi A, Persaud SJ: **A comparative analysis of human and mouse islet G-protein coupled receptor expression.** *Sci Rep* 2017, **7**:46600.
  59. Drynda R, Persaud SJ, Bowe JE, Jones PM: **the placental secretome: identifying potential cross-talk between placenta and islet beta-cells.** *Cell Physiol Biochem* 2018, **45**:1165-1171.
- This study identifies the GPCR ligands expressed by the placenta for which the cognate receptors are also expressed by the islets, identifying possible peptide signals that may potentially play a role in  $\beta$ -cell adaptation.
60. Dhillon WS, Savage P, Murphy KG, Chaudhri OB, Patterson M, Nijher GM, Foggo VM, Dancey GS, Mitchell H, Seckl MJ *et al.*: **Plasma kisspeptin is raised in patients with gestational trophoblastic neoplasia and falls during treatment.** *Am J Physiol Endocrinol Metab* 2006, **291**:E878-E884.
  61. Hauge-Evans AC, Richardson CC, Milne HM, Christie MR, Persaud SJ, Jones PM: **A role for kisspeptin in islet function.** *Diabetologia* 2006, **49**:2131-2135.
  62. Bowe JE, Foot VL, Amiel SA, Huang GC, Lamb M, Lakey J, Jones PM, Persaud SJ: **GPR54 peptide agonists stimulate insulin secretion from murine, porcine and human islets.** *Islets* 2012, **4**.
  63. Bowe JE, King AJ, Kinsey-Jones JS, Foot VL, Li XF, O'Byrne KT, Persaud SJ, Jones PM: **Kisspeptin stimulation of insulin secretion: mechanisms of action in mouse islets and rats.** *Diabetologia* 2009, **52**:855-862.
  64. Colledge WH: **Transgenic mouse models to study Gpr54/kisspeptin physiology.** *Peptides* 2009, **30**:34-41.
  65. Cetkovic A, Miljic D, Ljubic A, Patterson M, Ghatei M, Stamenkovic J, Nikolic-Djurovic M, Pekic S, Doknic M, Glisic A *et al.*: **Plasma kisspeptin levels in pregnancies with diabetes and hypertensive disease as a potential marker of placental dysfunction and adverse perinatal outcome.** *Endocr Res* 2012, **37**:78-88.
  66. Izzi-Engbeaya C, Comninou AN, Clarke SA, Jomard A, Yang L, Jones S, Abbara A, Narayanaswamy S, Eng PC, Papadopoulou D *et al.*: **The Effects of Kisspeptin on beta-cell Function, Serum Metabolites and Appetite in Humans.** *Diabetes Obes Metab* 2018.
- This study shows that kisspeptin administration in humans is able to safely and effectively stimulate insulin secretion, raising the possibility of future therapeutic use.
67. Butler AE, Cao-Minh L, Galasso R, Rizza RA, Corradin A, Cobelli C, Butler PC: **Adaptive changes in pancreatic beta cell fractional area and beta cell turnover in human pregnancy.** *Diabetologia* 2010, **53**:2167-2176.